



İZMİR MATHEMATICS DAYS - IV

JUNE 15-17, 2022

Izmir Institute of Technology

For registrations and all information

Visit: <https://img.iyte.edu.tr/>

Venue: Online

Registration Deadline

With presentation: 20 May 2022

Without presentation: 1 June 2022

Invited Speakers

İzzet Coşkun	University of Illinois at Chicago
Türkü Özlüm Çelik	Boğaziçi University
Burak Erdoğan	University of Illinois at Urbana-Champaign
Cihan Okay	Bilkent University
Sinem Onaran	Hacettepe University
Türker Özsarı	Bilkent University
Jan Stovicek	Charles University
Julian Weight	Aalto University
Joshua Zahl	University of British Columbia



İZMİR MATHEMATICS DAYS - IV

ABSTRACT BOOK

İZMİR INSTITUTE OF TECHNOLOGY
JUNE 15-17, 2022

Dear Participants,

“İzmir Mathematics Day” is an annual event organized by different universities in İzmir. It brings together distinguished mathematicians from our country and abroad with students and young researchers. One of the main goals of İzmir Mathematics Days is to create a platform where students and young researchers can share their studies, ideas and experiences, and can develop their research and mentor networks. Other aims are to introduce undergraduate and graduate students with new and interesting topics in advanced and modern mathematics and to encourage them to pursue a career as a researcher in mathematics or other related sciences.

The first İzmir Mathematics Days was hosted by Yaşar University in 2018, and the second took place at Dokuz Eylül University in 2019. Due to the COVID-19 pandemic in 2020, the third one was organized online by the mathematics department of Dokuz Eylül University. We appreciate the efforts of all our colleagues for establishing and organizing these previous meetings.

This year, we are happy to host the fourth İzmir Mathematics Days at İzmir Institute of Technology and say welcome to all participants. As members of the mathematics community in İzmir, we are proud to say that each year the interest to these days increases, with nearly 200 participants today and speakers from Turkey, USA, Canada, Finland and Czech republic.

The mathematics days this year are again organized to be online. Each day there will be a single morning session and there will be two parallel afternoon sessions. In the morning sessions, invited speakers will give colloquium talks introducing their fields of research. The afternoon sessions are reserved for the presentations given by graduate students and young researchers. Besides, on the evening of the first and second days, there will be two more invited talks. Most of the talks will be in English. In the present booklet, you can find all abstracts and program details of the event.

Here, we would like to thank all reputable, mathematicians İzzet Coşkun, Türkü Özlüm Çelik, Burak Erdoğan, Cihan Okay, Sinem Onaran, Türker Özsarı, Jan Stovicek, Julian Weight and Joshua Zahl for accepting our invitation. We believe their talks will enrich our research scope and inspire all young mathematicians.

Also, we would like to thank all students and young researchers who will contribute by their talks and participation. We hope this event will be a good opportunity to introduce your current work and communicate it with your peers and senior mathematicians. We believe you will enjoy and learn much during these days.

Finally, we would like to express our gratitude and appreciation to the Scientific Committee and the Academic Staff of the mathematics department at İzmir Institute of Technology, who contributed to the realization of the present fourth İzmir Mathematics Days.

Yours Sincerely, Organizing Committee

İzmir Institute of Technology, June 15-17, 2022

SCIENTIFIC COMMITTEE

Prof. Dr. Ergün Yalçın	<i>Bilkent University</i>
Prof. Dr. Halil Oruç	<i>Dokuz Eylül University</i>
Prof. Dr. Oktay Pashaev	<i>İzmir Institute of Technology</i>
Prof. Dr. Refail Alizade	<i>Yaşar University</i>
Prof. Dr. Selçuk Demir	<i>Dokuz Eylül University</i>
Asst. Prof. Dr. Asgar Jamneshan	<i>Koç University</i>
Asst. Prof. Dr. Aslı Güçlükan İlhan	<i>Dokuz Eylül University</i>

ORGANIZING COMMITTEE

Prof. Dr. Engin Büyükaşık	<i>İzmir Institute of Technology</i>
Prof. Dr. Oğuz Yılmaz	<i>İzmir Institute of Technology</i>
Prof. Dr. Şirin Atılğan Büyükaşık	<i>İzmir Institute of Technology</i>
Asst. Prof. Dr. Faruk Temur	<i>İzmir Institute of Technology</i>
Asst. Prof. Dr. Haydar Göral	<i>İzmir Institute of Technology</i>
Inst. Dr. Barış Çiçek	<i>İzmir Institute of Technology</i>
Res. Asst. Aylin Bozacı Serdal	<i>İzmir Institute of Technology</i>
Res. Asst. Aykut Alkın	<i>İzmir Institute of Technology</i>
Res. Asst. Hikmet Burak Özcan	<i>İzmir Institute of Technology</i>
Res. Asst. Kemal Cem Yılmaz	<i>İzmir Institute of Technology</i>
Res. Asst. Sinem Benli Göral	<i>İzmir Institute of Technology</i>

TABLE OF CONTENTS

Preface	i
Committees	
Scientific Committee	ii
Organizing Committee	ii
Table of Contents	iii
Program	vii
Invited Speakers	
<i>Burak Erdoğan</i>	
Dispersive Estimates for Dirac Operators	1
<i>Cihan Okay</i>	
Topology, Probability and Quantum	2
<i>İzzet Coşkun</i>	
Points in the Projective Plane	3
<i>Jan Stovicek</i>	
On the Finitistic Dimension Conjecture	4
<i>Joshua Zahl</i>	
The Kakeya Conjecture: A Survey	5
<i>Julian Weigt</i>	
Endpoint Regularity of Maximal Operators in Higher Dimensions	6
<i>Sinem Onaran</i>	
Open Book Decompositions, Contact 3-Manifolds and Legendrian Knots ...	7
<i>Türker Özşarı</i>	
Existence of Unattainable States for the Schrödinger Type Flows	8
<i>Türkü Özlüm Çelik</i>	
Algebraic Curves, Computer Algebra and Integrable Systems	9
Contributed Talks	
<i>Abdubaki Aşur</i>	
John's Theorem and Series in Banach Spaces	10
<i>Aygül Koçak Özvarol</i>	
Post Quantum PQ Calculus and Supersymmetric Units of Quantum Information	11
<i>Aykut Alkın</i>	
The Initial-Boundary Value Problem for Higher Order Schrödinger Equation on the Half-Line	12
<i>Aylin Bozacı Serdal</i>	
Exactly Solvable Generalized Burgers Models with Variable Coefficients ..	13

<i>Ayşe Fidan</i>	
Nonexistence of Solutions for the Variable Coefficients Wave Equation with Strong Damping	14
<i>Badik Hüseyin Uysal</i>	
Some Geometric Properties of Orlicz Spaces Equipped with s-norms	15
<i>Cihat Aydın</i>	
Relationship of Fixed Point Theory with Gradient Descent Methods	16
<i>Çağatay Altuntaş</i>	
Dedekind Harmonic Numbers	17
<i>Derya Özdemir</i>	
Qualitative Properties of Solutions of Some Keller - Segel Type Systems ..	18
<i>Ege Tamcı</i>	
Stability Analysis and Control of the Kuramoto Model Driven with Alpha Stable Processes	19
<i>Engin Başakoğlu</i>	
Smoothing and Global Attractors for the Hirota-Satsuma System on the Torus	20
<i>Eren Şen</i>	
Trisections of Smooth 4-Manifolds	21
<i>Esma Dirican Erdal</i>	
A Formula for Volume forms on Representation Varieties of Surfaces with Boundary $\Sigma_{g,2}$	22
<i>Esra Başar</i>	
A Full Description of Compact Extremal Operators on Orlicz Spaces Equipped with s-norms	23
<i>Ezgi Gürbüz</i>	
Unique Decomposition into Regular Ideals for Marot Rings	24
<i>Gamze Akar</i>	
Rational Groups whose Character Degree Graphs are Disconnected	25
<i>Gürcihan Zaman</i>	
Mathematical Modeling of Infectious Diseases	26
<i>Haleh Hamdi</i>	
Bazzoni's Conjecture and Almost Prüfer Domains	27
<i>Hazal Yüksekaya</i>	
Mathematical Behavior of Solutions for a Delayed Hyperbolic-Type Equations	28
<i>İlayda Kaplan</i>	
On 1-Absorbing Intuitionistic Fuzzy Ideals of Commutative Semirings	29
<i>İdem Susuzlu</i>	
Existence and Stabilization of Solutions of a Linear Viscoelastic Wave Equation with Neumann Boundary Condition	30

<i>Kemal Cem Yılmaz</i>	
Stabilization of Higher-Order Schrödinger Equation Using Right Endpoint Controllers	31
<i>Kübra Akyıl</i>	
3 Noktalı Yol Grafin Karışık Genişlemelerinin Spektral Karakterizasyonu .	32
<i>Kübra Karaağaç</i>	
Metrik Uzaylar Üzerinde Bazı Sabit Çember Teoremleri ve Örnekleri	33
<i>Meliha Akcan</i>	
İşaretli Grafların Spektral Özellikleri	34
<i>Merve Bülbül</i>	
Sıra Norm Sürekli Operatörlerin Latis Yapısı Üzerine	35
<i>Mustafa Kutay Kutlu</i>	
Roth's Theorem on Arithmetic Progressions	36
<i>Oğuz Şavk</i>	
New Infinite Rank Summands of Homology Cobordism Group	37
<i>Osman Hamza</i>	
Truncation on ℓ -groups	38
<i>Özlem Moralı Kızanlık</i>	
Metrik Uzaylarda Sabit Çemberler	39
<i>Pınar Şaşmaz</i>	
ωe^* -Ayrırma Aksiyomları	40
<i>Rabia Ayan</i>	
On 2-Absorbing Vague Fuzzy Ideals of Commutative Semirings	42
<i>Sabahat Defne Plattürk</i>	
Dirichlet Boundary Value Problem for the Laplace Equation in the Semi-Infinite Domain via a Boundary Integral Equation Method	43
<i>Sinem Benli Göral</i>	
A Generalization of Injective Modules	44
<i>Şeyma Yaşar</i>	
Classification of Orlicz Spaces Equipped with s-norms	45
<i>Şule Kılıçaslan</i>	
Spin Structures on Real Bott Manifolds	46
<i>Tuğrul Cömert</i>	
Asymptotic Behavior of Solutions for p-biharmonic Parabolic Equation with Logarithmic Nonlinearity	47
<i>Uyen Enni Le</i>	
On the Depth and Reflexivity of Modules	48
<i>Victor Blasco</i>	
Categorical Methods in the Representation Theory of Artin Algebras	49
<i>Yeşim Demiroğlu Karabulut</i>	
Applications of Cayley Digraphs to Waring's Problem and Sum-Product Formulas	50

<i>Yonca Ünver</i>	
Döngüsel Çizgelerin Ricci Eğrilikleri	51
<i>Zehra Çayıç</i>	
Time-evolution of Cauchy-Euler Type Quantum Oscillator in the Presence of Variable Magnetic and Electric Fields	52
<i>Zeynep Kara</i>	
İki Değişkenli q -Bleimann, Butzer ve Hahn Operatörlerinin Yaklaşım Oranları	53
List of Participants	54

PROGRAM

		15-Jun-22	
9:45	10:00	Opening	
10:00	11:00	Invited Speaker : BURAK ERDOĞAN	
11:00	11:15	Break	
11:15	12:15	Invited Speaker : SİNEM ONARAN	
12:15	13:30	Lunch Break	
13:30	14:30	Invited Speaker : CİHAN OKAY	
14:30	14:35	Break	
		İMG 2022 - SECTION 1	İMG 2022 - SECTION 2
14:35	15:05	Abdulkaki Aşur	Oğuz Şavk
15:05	15:35	Özlem Moral Kızanlık	Esmâ Dirican Erdal
15:35	16:05	Esra Başar	Pınar Şaşmaz
16:05	16:15	Break	
16:15	16:45	Şeyma Yaşar	Yonca Ünver
16:45	17:15	Badik Hüseyin Uysal	Şule Kılıçaslan
17:15	17:45	Merve Bülbül	Eren Şen
17:45	18:30	Break	
18:30	19:30	Invited Speaker : JOSHUA ZAHL	

		16-Jun-22	
9:30	10:30	Invited Speaker : TÜRKÜ ÖZLÜM ÇELİK	
10:30	10:45	Break	
10:45	11:45	Invited Speaker : JAN STOVICEK	
11:45	11:50	Break	
		İMG 2022 - SECTION 1	İMG 2022 - SECTION 2
11:50	12:20	Yeşim Demiroğlu Karabulut	Hazal Yüksekaya
12:20	13:30	Lunch Break	
13:30	14:00	Zeynep Kara	Aykut Alkın
14:00	14:30	Meliha Akcan	Kemal Cem Yılmaz
14:30	15:00	Haleh Hamdi	Tuğrul Cömert
15:00	15:15	Break	
15:15	15:45	Gamze Akar	Zehra Çayıç
15:45	16:15	Ezgi Gürbüz	Gürcihan Zaman
16:15	16:45	Çağatay Altuntaş	Ege Tamcı
16:45	17:00	Break	
17:00	17:30	Kübra Karaağaç	İdem Susuzlu
17:30	18:00	Break	
18:00	19:00	Invited Speaker : İZZET COŞKUN	

		17-Jun-22	
9:30	10:30	Invited Speaker : TRKER ZSARI	
10:30	10:45	Break	
10:45	11:45	Invited Speaker : JULIAN WEIGT	
11:45	11:50	Break	
		İMG 2022 - SECTION 1	İMG 2022 - SECTION 2
11:50	12:20	Kbra Akyıl	Engin Bařakoęlu
12:20	13:30	Lunch Break	
13:30	14:00	Osman Hamza	Aylin Bozacı Serdal
14:00	14:30	Mustafa Kutay Kutlu	Cihat Aydın
14:30	15:00	Sinem Benli Gral	Sabahat Defne Plattrk
15:00	15:15	Break	
15:15	15:45	Victor Blasco	Aygl Koęak zvarol
15:45	16:15	İlayda Kaplan	Ayře Fidan
16:15	16:45	Rabia Ayan	Derya zdemir
16:45	17:15	Uyen Le	
17:15	17:30	Closing	

Dispersive Estimates for Dirac Operators

Burak Erdoğan

University of Illinois at Urbana-Champaign

`berdogan@illinois.edu`

We will review recent results on dispersive decay estimates for the Dirac operators in \mathbb{R}^n with a decaying self-adjoint potential. These include $L^1 \rightarrow L^\infty$ and Strichartz estimates. We also discuss the classification of threshold resonances and their effects on dispersive estimates and the limiting absorption principle for Dirac operators. The talk is based on joint works with M. Goldberg (U. Cincinnati), W. Green (RoseHulman), and E. Toprak (Yale).

Topology, Probability and Quantum

Cihan Okay

Bilkent University

`cihan.okay@bilkent.edu.tr`

Our experience of the physical world is essentially shaped by the act of measurement. In abstract terms a measurement can be described by a probability table consisting of a collection of measurements and the probabilities assigned to the occurrence of each outcome. When there are more than two parties, say Alice and Bob, performing measurements the resulting probability tables are dramatically different for quantum measurements, than their classical counterparts. The mathematical theory of quantum mechanics is based on Hilbert spaces and operators acting on them and Born's rule tells us how to extract probabilities as the result of a measurement. A quirky feature of quantum is observed when Alice and Bob perform measurements on entangled states. In these cases the probability tables cannot be reproduced by classical measurements. This feature, known as quantum contextuality, plays a prominent role in computational advantage in quantum computers. Topology enters the picture at this point as an intrinsic structure of the collection of measurements. In this talk I will explain how the theory of simplicial sets, combinatorial representations of topological spaces in modern homotopy theory, can be enriched into a probabilistic version that can be used to study this fundamental quantum phenomenon.

Points in the Projective Plane

İzzet Coşkun

University of Illinois at Chicago

`coskun@math.uic.edu`

In this talk, I will discuss the geometry of the Hilbert scheme of n points in the projective plane, which is a smooth compactification of the configuration space of n points. I will focus on the question: What is the most special codimension one position that n points can lie in? For example, three points are typically not collinear, but in codimension one they can be collinear. This simple question will lead us to a tour of some fun mathematics ranging from moduli spaces of stable sheaves on the plane to fractal curves and palindromic numbers. This talk is based on joint work with Jack Huizenga and Matthew Woolf.

On the Finitistic Dimension Conjecture

Jan Stovicek

Charles University

`stovicek@karlin.mff.cuni.cz`

Given a ring, a basic question is what the modules (also known as representations) over the ring look like. Here we restrict to the cases of (1) commutative noetherian rings (typical examples are coordinate rings of algebraic varieties or the rings of integers of number fields) and of (2) finite dimensional algebras over a field. Homological algebra provides a powerful toolbox for studying the structure of modules and it is a basic question what homological dimensions modules can attain. The Finitistic Dimension Conjecture, going back to the 1960's, says that for each ring belonging to one of the two classes mentioned above, the projective dimension of a module can only have finitely many values. Whereas in the commutative noetherian case the problem is well understood and the conjecture is false, in the situation of finite dimensional algebras the problem resists for more than half a century despite numerous efforts to solve it. In the first part of the talk, I will explain the conjecture along with some important results, while in the second part I will focus on recent contributions.

The Kakeya Conjecture: A Survey

Joshua Zahl

University of British Columbia
`jzahl@math.ubc.ca`

The Kakeya conjecture is a problem in geometric measure theory that has connections to several deep open problems in harmonic analysis. In this talk, I will discuss the history of the Kakeya conjecture, and some of the ideas that have been used to make progress on the problem in recent years.

Endpoint Regularity of Maximal Operators in Higher Dimensions

Julian Weigt

Aalto University
julian.weigt@aalto.fi

We prove the endpoint regularity bound

$$\|\nabla Mf\|_{L^1(\mathbb{R}^n)} \leq C_n \cdot \|\nabla f\|_{L^1(\mathbb{R}^n)}$$

for some maximal functions Mf in any dimension $n \geq 1$. This bound and various related questions have been intensively studied for all sorts of maximal operators. Nevertheless, it has essentially only been resolved on L^p for $p > 1$ and in one dimension. We prove it for the uncentered Hardy-Littlewood maximal function of characteristic functions, for general dyadic maximal functions and for the cube maximal function.

The key arguments of the proofs are of geometric nature. For example new variants of the isoperimetric inequality and of the Vitali Covering Lemma are proven and used. All proofs are mostly elementary up to applications of classical results like the relative isoperimetric inequality and the coarea formula.

Open Book Decomposition, Contact 3-Manifolds and Legendrian Knots

Sinem Onaran

Hacettepe University
`sonaran@hacettepe.edu.tr`

Due to Alexander, it is well known that every closed oriented 3-manifold has an open book decomposition. In the first part of the talk, I will define open book decompositions of 3-manifolds and discuss various examples in detail. Further, we I will discuss the importance of the open books in manifold theory, in particular in contact geometry. A contact structure on an odd-dimensional manifold is a maximally nonintegrable hyperplane field that vanishes nowhere. After a brief introduction to contact 3-manifolds, I will focus on a class of knots called Legendrian knots in contact 3-manifolds. A knot in a contact 3-manifold is called Legendrian if it is always tangent to the contact planes. I will discuss a new invariant for Legendrian knots which is defined using open book decompositions and discuss its applications.

Existence of Unattainable States for the Schrödinger Type Flows

Türker Özsari

Bilkent University

`turker.ozsari@bilkent.edu.tr`

I will first introduce the notion of controllability for differential equations in certain senses starting with the finite dimensional case. Then, I will move to distributed parameter systems (PDEs) and I will contrast the two cases. Using an integral transform method, I will give an elementary proof of the existence of physically reasonable states to which the heat flow cannot be steered. I will show that the same technique fails for dispersive equations which will motivate us to consider the latter problem with rather more technical tools. These include recent developments in well-posedness theory of initial- /boundary value problems as well as control theoretical techniques such as the Hilbert Uniqueness Method. We will perform this with the canonical example of the classical Schrödinger equation and also extend the same results to biharmonic problems.

Algebraic Curves, Computer Algebra and Integrable Systems

Türkü Özlüm Çelik

Boğaziçi University
turkuozlum@gmail.com

An important application of algebraic curves lies in the context of integrable systems via theta functions. They give rise, for instance, to solutions of the Kadomtsev-Petviashvili hierarchy, a universal integrable system. This talk aims to make an excursion into this study from the lens of computational algebraic geometry. The emphasis will be on exploiting modern tools in symbolic, numerical and combinatorial algebraic geometry to investigate such solutions and their applications back to the algebraic curves.

John's Theorem and Series in Banach Spaces

Abdurbaki Aşur

Dokuz Eylül University
abdurbaki.asur@ogr.deu.edu.tr

Co-author: Selçuk Demir
selcuk.demir@deu.edu.tr

We are going to present a proof of a theorem of F. John characterizing ellipsoids of maximal volume in symmetric convex bodies in \mathbb{R}^n . The proof which will be presented is a relatively recent one due to F. Schuster and P.M. Gruber. We also present some basic results in Banach Space Theory which are related somehow to the geometric ideas present in John's theorem.

References

- [1] F. Schuster, P.M. Gruber. An Arithmetic Proof of John's Theorem, *Archiv der Mathematik*, **85**, 82-88 (2005).
- [2] Extremum problems with inequalities as subsidiary conditions, in: *Studies and essays presented to R. Courant on his 60th birthday*, January 8, 1948, 187-204, Inter-science, New York 1948.
- [3] A. Dvoretzky, C.A. Rogers. Absolute and Unconditional Convergence in Normed Linear Spaces, *PNAS 1950*; 36, 192-197

Post Quantum PQ Calculus and Supersymmetric Units of Quantum Information

Aygül Koçak Özvarol

İzmir Institute of Technology
aygulkocak@iyte.edu.tr

Co-author: Oktay Pashaev
oktaypashaev@iyte.edu.tr

In this talk first we introduce main ingredients of Quantum Calculus with two arbitrary bases P and Q , which is called also as Post Quantum Calculus of finite differences [1]. The calculus is related with representation of quantum groups and includes as particular cases the non-symmetric q -calculus, symmetric q -calculus, Fibonacci-Golden calculus [2], calculus of Fibonacci divisors [3] and Tamm-Dankoff quantum calculus. By introducing PQ-oscillator algebra and corresponding PQ-coherent states, we describe Fock-Bargmann representation of quantum states in terms of this calculus. In the framework of supersymmetric quantum mechanics, the generalized PQ-supersymmetric annihilation operator and corresponding eigenstates as supersymmetric PQ-coherent states are derived. These states reflect hidden PQ-Quantum supersymmetry and can be expanded in terms of orthonormal fermionic and bosonic coherent states. We show that these coherent states can represent unit of quantum information, parametrized by points on Supersymmetric Bloch Sphere, where bosonic and fermionic states correspond to the North and the South poles of the sphere, correspondingly. The Shannon entropy of our states takes maximal value for the maximally random super-qubit states on the Equator circle of Supersymmetric Bloch Sphere.

This work was supported by BAP project 2022 IYTE-1-0002.

Keywords. q -calculus, supersymmetry, super-qubit, coherent states.

References

- [1] Pashaev, O.K. PQ-Calculus of Fibonacci Divisors and Method of Images in Planar Hydrodynamics, *Mathematical Methods For Engineering Applications: ICMASE 2021, Springer Proceeding in Mathematics&Statistics* **384**(125-136) (2022).
- [2] Pashaev, O.K and Nalci, S. Golden quantum oscillator and Binet-Fibonacci calculus, *J. Phys. A: Math. Theor.* **45**(015303) (2012).
- [3] Pashaev, O.K. Quantum calculus of Fibonacci divisors and infinite hierarchy of Bosonic-Fermionic quantum oscillators, *Int. J. Geom. Methods in Modern Physics* **18**(2150075) (2021).

The Initial-Boundary Value Problem for Higher Order Schrödinger Equation on The Half-Line

Aykut Alkın

İzmir Institute of Technology
aykutalkin@iyte.edu.tr

Co-author: Türker Özşarı
turker.ozsari@bilkent.edu.tr

The initial-boundary problem for higher order Schrödinger equation on the right half-line is considered with a source f , an initial value u_0 and an inhomogeneous Dirichlet datum g in L^2 -based fractional Sobolev spaces:

$$\begin{aligned} iu_t + i\beta u_{xxx} + \alpha u_{xx} + i\delta u_x &= f, & (x, t) \in \mathbb{R}_+ \times (0, T), \\ u(x, 0) &= u_0(x), & x \in \mathbb{R}_+, \\ u(0, t) &= g(t), & t \in (0, T), \end{aligned} \tag{1}$$

where $\alpha, \delta \in \mathbb{R}$ and $T, \beta > 0$. We analyze the regularity of the solution $u(x, t)$ of the linear ibvp (1) by using a decompose-reunify argument. We first decompose the linear ibvp into the pure homogeneous ibvp with zero initial datum, the homogeneous ivp and the nonhomogeneous ivp with zero initial datum to observe separately the effect of each data f, u_0 and g on the regularity. After estimating the solutions of the decomposed problems with respect to their own inhomogeneous data, respectively, we reunify them stating the relation between the data (f, u_0, g) for the linear ibvp (1) and the corresponding data for the decomposed problems. We also obtain a representation formula for the pure homogeneous ibvp with zero initial datum by using the unified transform method, which is also known as Fokas' method [1]. Consequently, this formulation is used to obtain space and time estimates for the corresponding model.

This study is a part of our local and global well-posedness analysis for higher order nonlinear Schrödinger equation.

Keywords. Fokas' method. Space-time estimates. Local wellposedness. Global well-posedness. Regularity.

References

- [1] A. S. Fokas, A unified approach to boundary value problems, *Vol 78 of CBMS-NSF Regional Conference Series in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA* (2008).

Exactly Solvable Generalized Burgers Models with Variable Coefficients

Aylin Bozacı Serdal

İzmir Institute of Technology
aylinbozaci@iyte.edu.tr

Co-author: Şirin Atılğan Büyükaşık
sirinatilgan@iyte.edu.tr

In this work, we study an initial value problem (IVP) for a one dimensional generalized Burgers equation (BE) with variable coefficients, $U_t + (\dot{\mu}(t)/\mu(t))U + UU_x = (1/2\mu(t))U_{xx} - a(t)U_x + b(t)(xU)_x - \omega^2(t)x + f(t)$, defined for $x \in \mathbb{R}$, $t > 0$, [1],[2]. First, by using Cole-Hopf transformation, we linearize the generalized BE to the diffusion-convection-reaction equation and then applying Wei-Norman Lie algebraic procedure [1], we find the evolution operator of the corresponding linear diffusion-type equation exactly in terms of solutions to the second order inhomogeneous characteristic ODE determined by the time-dependent parameters of the generalized BE. Then, we formulate the general closed form solution of the IVP for generalized BE in terms of this characteristic equation and solutions of IVP for standard BE. Later, using the translation and Galilean symmetries of the standard BE, we construct families of Burgers solutions whose wave profiles propagate according to the general solution of the characteristic equation. Also we discuss some particular solutions such as generalized shocks, triangular and N-shaped generalized traveling waves and rational type solutions with pole type moving singularities and introduce concrete exactly solvable models, [4].

Keywords. Generalized Burgers equation, variable coefficients, initial value problem, evolution operator method, exact solutions.

Reference

- [1] J.M. Burgers, A Mathematical Model Illustrating the Theory of Turbulence. *Adv. Appl. Mech.*, **1**,171, (1948).
- [2] Axel Schulze-Halberg, Burgers equation with time-dependent coefficients and nonlinear forcing term: Linearization and exact solvability, *Commun Nonlinear Sci Numer Simulat.* **22**, 1068-1083,(2015).
- [3] J. Wei, E. Norman, Lie algebraic solution of linear differential equations, *J.Math.Phys.* **4**: 575, (1963).
- [4] Ş.A. Büyükaşık, A. Bozacı, Dynamical properties of generalized traveling waves of exactly solvable forced Burgers equations with variable coefficients. *Commun Nonlinear Sci Numer Simulat.* **96**, (2021), 105682.

Nonexistence of Solutions for the Variable Coefficients Wave Equation with Strong Damping

Ayşe Fidan

Dicle University
afidanmat@gmail.com

Co-author: Erhan Pişkin
episkin@dicle.edu.tr

In this work, we consider the variable coefficients wave equation with strong damping. Under suitable conditions on variable coefficients, we prove the blow up of solutions.

Keywords. Nonexistence, Wave equation, Variable coefficients.

References

- [1] V. Georgiev, G. Todorova. Existence of a solution of the wave equation with nonlinear damping and source terms, *Journal of Differential Equations* **109**(2)(1994) 295-308.
- [2] E. Pişkin, B. Okutmuşur. An Introduction to Sobolev Spaces, *Bentham Science* (2021).
- [3] X. Zheng, Y. Shang, X. Peng. Blow up of Solutions for a Nonlinear Petrovsky Type Equation with Time-dependent Coefficients, *Acta Mathematicae Applicatae Sinica, English Serie* **36**(4)(2020) 836-846.

Some Geometric Properties of Orlicz Spaces Equipped with s -norms

Badik Hüseyin Uysal

İstanbul University
huseyinuyisal@istanbul.edu.tr

Co-author: Serap Öztop Kaptanoğlu, Esra Başar, Şeyma Yaşar
oztops@istanbul.edu.tr, esra.basar@yeditepe.edu.tr, seymayasar@gtu.edu.tr

Let (X, Σ, μ) σ -finite, non-atomic and complete measurable space and Φ be an Orlicz function. In Orlicz space L^Φ , the first aim is to give a relation the other norms with s -norm family on Orlicz spaces. The second aim is to examine some geometric properties in Orlicz spaces with respect to s -norms, which are studied with respect to Luxemburg, Orlicz and Amemiya norms. In this study, some geometric properties of Orlicz space are given in simpler terms.

Keywords. Orlicz Space, Extreme Points, Compact Operators.

References

- [1] Chen, S. and Wisła, M., Extreme compact operators from Orlicz spaces to $C(\Omega)$, *Comment. Math. Univ. Carolin.*, 34(1), (1993)
- [2] Dunford, N. and Schwartz, J.T., *Linear Operators I: General Theory*, Interscience, (1958).
- [3] M. Wisła, Orlicz spaces equipped with s -norms, *Journal of Mathematical Analysis and Applications*, **483**(2), (2020).
- [4] Rao, M.M. and Ren, Z.D., *Theory of Orlicz Spaces*, Marcel Dekker, (1991).

Relationship of Fixed Point Theory with Gradient Descent Methods

Cihat Aydın

Yıldız Technical University
cihat.aydin@std.yildiz.edu.tr

It is well known that gradient descent methods can be realized by using some well known fixed point iterations. In fact, the main purpose of gradient descent methods is to minimize the objective functions associated with a concerned problem by using these iterations. Especially, optimization techniques based on gradient methods are of the utmost importance in minimizing error functions used to train a machine(or deep) learning algorithm. In particular, when the size of data is sufficiently large, gradient descent methods requires to be accelerated [1]. To manage this acceleration task, many works were brought to the literature using fixed point iterations [1],[2]. In this presentation, we first mention about gradient descent methods and their accelerations.

Keywords. convex optimization, fixed point theory, big data, machine learning, contraction mapping, gradient descent, heavy balls.

References

- [1] A. Jung. A Fixed-Point of View on Gradient Methods for Big Data, *Frontiers in Applied Mathematics and Statistics* **3**(18) (2017).
- [2] H. F. Walker and P. Ni. Anderson acceleration for fixed-point iterations, *SIAM J. Numer. Anal.* **49**(4) (2011).

Dedekind Harmonic Numbers

Çağatay Altuntaş

İstanbul Technical University
caltuntas@itu.edu.tr

We introduce a generalization of the harmonic numbers, motivated by the Dedekind zeta function of a number field. For any number field K , we define $h_K(n)$, the Dedekind harmonic number, as the sum of reciprocals of the norm of integral ideals with norm less than n . We will show that for any number field, the corresponding Dedekind harmonic numbers are non-integer except for finitely many of them, together with density results.

Keywords. harmonic numbers, prime number theory, Dedekind zeta function, number fields.

References

- [1] Ç. Altuntaş, H. Göral, (2021). Dedekind harmonic numbers. *Proc. Math. Sci.* 131(46).

Qualitative Properties of Solutions of Some Keller - Segel Type Systems

Derya Özdemir

İzmir Institute of Technology
deryacokgur@iyte.edu.tr

Co-author: Assoc. Prof. Dr. Türker Özsarı
turker.ozsari@bilkent.edu.tr

Mathematical modeling of biological phenomena has become important to better understand the associated process. One of the these biological events is chemotaxis. Keller - Segel type system first presented in 1970 is a classical chemotaxis modeling [1]. This model includes a system of partial differential equations. I will introduce some Keller - Segel type systems and talk about qualitative properties of their solutions.

Keywords. . Chemotaxis, local existence, global existence, boundedness, blow-up, Keller - Segel model

References

- [1] E. F. Keller and L. A. Segel . Initiation of slime mold aggregation viewed as an instability, *Journal of Theoretical Biology* **26**(3) (1970).

Stability Analysis and Control of the Kuramoto Model Driven with Alpha Stable Processes

Ege Tamcı

İzmir Institute of Technology
egetamci@iyte.edu.tr

Co-authors:

Ferit Acar Savacı¹, Faruk Temur²

¹acarsavaci@iyte.edu.tr, ²faruktemur@iyte.edu.tr

Finite variance Gaussian processes are insufficient in modelling the noise with irregular and impulsive behavior in the power grids modeled with Kuramoto Model with alpha-stable process. For this reason, using the alpha-stable process (alpha(α)-stable Levy process), which have infinite variance and heavy tailed distributions will be more appropriate model for the noise in the system. Although alpha stable processes better model the noise in the system, it is very difficult to analyze the stability of the system using the Lyapunov theory. In this study, we indicate how we can take advantage of the Lyapunov theory by making use of the integral representation with respect to a Poisson random measure of alpha stable processes. Thus, we obtain a controller for the stochastic Kuramoto model driven alpha stable processes and determine the conditions for the parameters in the model to provide the stochastic stability in the asymptotic sense.

Keywords. Non-Gaussian processes; Non-linear processes; Stochastic control; Power Systems

Smoothing and Global Attractors for the Hirota-Satsuma System on the Torus

Engin Başakoğlu

Boğaziçi University

`engin.basakoglu@boun.edu.tr`

In this talk, we consider the Hirota-Satsuma system, a coupled Korteweg de Vries system (CKdV), with periodic boundary conditions. The model explains two long waves' interactions with separate dispersion relations. CKdV systems have a wide range implementations in different fields of chemistry, biology, hydrodynamics, mechanics, plasma physics, etc. The talk concerns with the dynamics of solutions to the Hirota-Satsuma system. In particular, given initial data in a Sobolev space, the difference of the nonlinear and linear evolutions lies in a smoother space. The smoothing gain we obtain depends very much on the arithmetic nature of the coupling parameter of the system. To cope with the resonances stemming from interactions of various frequencies, dependent on the coupling parameter, we invoke the Diophantine approximation of real numbers. Moreover, we address the forced and damped version of the Hirota-Satsuma system, once having the analogous smoothing estimates, we discuss the existence and smoothness of a global attractor in the energy space.

Trisections of Smooth 4-Manifolds

Eren Şen

Middle East Technical University
`sen.eren@metu.edu.tr`

Trisections of 4-manifolds were first introduced by David Gay and Robion Kirby [1] and have blossomed into a popular field of study ever since. As trisections are in close relation with Heegaard splittings of 3-manifolds, they carry three-dimensional techniques into the strange realm of dimension four.

In our talk, we will state some well-known results and open problems from the trisections literature after introducing essential parts of the trisections theory including stabilizations, trisection diagrams, and the relation with handle decompositions. When a 4-manifold has connected boundary, we will see that a trisection yields an extra structure [1, 3]. Toward the end of the talk, we will give algorithms for passing between Kirby diagrams and trisection diagrams [2, 3], and describe a new relative trisection diagram for a Mazur Manifold.

Keywords. 4-manifold, Heegaard splitting, trisection, Mazur manifold.

References

- [1] David T. Gay and Robion Kirby. Trisections of 4-manifolds, *Geometry & Topology* **20** 3097-3132 (2016).
- [2] Nickolas A. Castro, David T. Gay, and Juanita Pinzón-Caicedo. Trisections of 4-manifolds with boundary. *PNAS* **115**(43) 10861-10868 (2018).
- [3] ———. Diagrams for Relative Trisections. *Pacific Journal of Mathematics* **294**(2) 275-305 (2018).

A Formula for Volume forms on Representation Varieties of Surfaces with Boundary $\Sigma_{g,2}$

Esma Dirican Erdal

Yeditepe University
esmadirican131@gmail.com.tr

Let $\Sigma_{g,n}$ be a compact oriented connected surface of genus $g = 2k \geq 4$ with boundary disjoint union of n circles. Let G be the complex reductive group $SL_2(\mathbb{C})$. We restrict our attention to irreducible representations whose stabilizers coincide with the center of G . In [1], such representations are called *good representations*. For surfaces $\Sigma_{g,n}$, we denote the set of conjugacy classes of good representations from the fundamental group of $\Sigma_{g,n}$ to G by $\mathcal{R}^g(\pi_1(\Sigma_{g,n}), G)$. The relative set of conjugacy classes of representations is given by $\mathcal{R}(\pi_1(\Sigma_{g,n}), \partial(\Sigma_{g,n}), G)_{\rho_0}$ and we denote the corresponding open subset of good representations by $\mathcal{R}^g(\pi_1(\Sigma_{g,n}), \partial(\Sigma_{g,n}), G)_{\rho_0}$ for a representation $\rho_0 : \pi_1(\Sigma_{g,n}) \rightarrow G$. The twisted Reidemeister torsion of $\Sigma_{g,n}$ can be seen as a holomorphic volume form $\Omega_{\Sigma_{g,n}}$ on $\mathcal{R}^g(\pi_1(\Sigma_{g,n}), G)$ and the holomorphic symplectic form $\omega_{\Sigma_{g,n}}$ is defined on relative representation variety $\mathcal{R}^g(\pi_1(\Sigma_{g,n}), \partial(\Sigma_{g,n}), G)_{\rho_0}$. In order to obtain a holomorphic symplectic volume form on $\mathcal{R}(\partial(\Sigma_{g,n}), G)$, Heusener and Porti identified the representation variety of the circle \mathbb{S}^1 with G , by mapping each representation to the image of a fixed generator of $\pi_1(\mathbb{S}^1)$. They restricted to representations which map the generators of $\pi_1(\mathbb{S}^1)$ to regular elements and established a relation between the volume forms $\Omega_{\Sigma_{g,n}}$ and $\omega_{\Sigma_{g,n}}$ [2]. By using this relation with certain conditions, we show that the holomorphic volume form on the representation variety of the surface $\Sigma_{g,2}$ can be expressed as a product of the symplectic holomorphic volume forms on the representation variety of $\Sigma_{2,2}$. Precisely,

$$\left| \Omega_{\Sigma_{g,2}}(\wedge \mathbf{h}_{\Sigma_{g,2}}^1) \right| = \mathcal{M}_k \prod_{i=1}^k \left| \omega_{\Sigma_{2,2}}^5(\wedge \mathbf{h}_{\Sigma_{2,2}}^1) \wedge \nu_1^* \wedge \nu_2^* \right|.$$

Here, $\mathcal{M}_k = (5!)^{-k} \frac{\nu(\wedge \mathbf{h}_{\mathbb{S}^1}^1)^{-2k+2}}{\langle \wedge \mathbf{h}_{\mathbb{S}^1}^1, \wedge \mathbf{h}_{\mathbb{S}^1}^0 \rangle^{-2k+2}} \in \mathbb{C}$.

Keywords. Twisted Reidemeister torsion, volume form, representation variety.

References

- [1] Johnson D. and Millson J., Deformation Spaces Associated to Compact Hyperbolic Manifolds, *Bull. Amer. Math. Soc.*, **14** (1986) 99-102.
- [2] Heusener M. and Porti J., Holomorphic volume forms on representation varieties of surfaces with boundary, *Annales Henri Lebesgue* **3** (2020) 341–380.

A Full Description of Compact Extremal Operators on Orlicz Spaces Equipped with s -norms

Esra Başar

Yeditepe University
esra.basar@yeditepe.edu.tr

Co-author:

Serap Öztop Kaptanoğlu, Badik Hüseyin Uysal, Seyma Yaşar
oztops@istanbul.edu.tr, huseyinuyisal@istanbul.edu.tr, seymayasar@gtu.edu.tr

Let Ω be a compact Hausdorff space, E be the predual of the Orlicz space L^Φ and T be a compact operator from E to $C(\Omega)$. The map $T' : \Omega \rightarrow L^\Phi$ is defined by $(T'\omega)(x) = (Tx)(\omega)$ for $\omega \in \Omega, x \in E$. Then T is called as nice if $T'(\Omega) \subseteq \text{Ext } B(L^\Phi)$ where $\text{Ext } B(L^\Phi)$ is the set of extreme points of closed unit ball in L^Φ . It is known that nice operators are extremal [2]. For the converse, under some topological properties of the set $\text{Ext } B(L^\Phi)$, we have that all compact extremal operators consist of nice operators. In this study, some of these properties for the set of extreme points in Orlicz spaces were investigated with respect to s -norms.

Keywords. Orlicz Spaces, Extreme Points, Compact Operators

References

- [1] Chen, S. and Wisła, M., Extreme compact operators from Orlicz spaces to $C(\Omega)$, *Comment. Math. Univ. Carolin.*, 34(1), (1993)
- [2] Dunford, N. and Schwartz, J.T., *Linear Operators I: General Theory*, Interscience, (1958).
- [3] M. Wisła, Orlicz spaces equipped with s -norms, *Journal of Mathematical Analysis and Applications*, **483**(2), (2020)
- [4] Rao, M.M. and Ren, Z.D., *Theory of Orlicz Spaces*, Marcel Dekker, (1991).

Unique Decomposition into Regular Ideals for Marot Rings

Ezgi Gürbüz

İzmir Institute of Technology
ezgigurbuz@iyte.edu.tr

Co-author: Başak Ay Saylam
basakay@iyte.edu.tr

Let R be a commutative ring with unity. In the papers [1], [2], and [3], the authors define the UDI property as: R has the unique decomposition into ideals (UDI) property if, for any R -module which decomposes into a finite direct sum of ideals, this decomposition is unique up to the order and isomorphism class of the ideals. They gave a characterization of a Noetherian domain/ring to satisfy UDI property.

We define the UDI-like property called UDRI property in the following way: R has the unique decomposition into regular ideals (UDRI) property if, for any R -module which decomposes into a finite direct sum of regular ideals, this decomposition is unique up to the order and isomorphism class of the regular ideals [4].

In this talk, we will present some preliminary results for Marot rings (every regular ideal of R is generated by its regular elements) whose regular ideals are finitely generated and give a necessary and sufficient condition for these rings to satisfy the UDRI property.

Keywords. Marot rings, valuation rings, h -local rings, regular localization

References

- [1] B. Ay, L. Klingler, Unique decomposition into ideals for reduced commutative Noetherian rings, *Trans. Am. Math. Soc.* **363** (7) (2011) 3703-3716.
- [2] P. Goeters and B. Olberding, Unique decomposition into Ideals for Noetherian domains, *Journal of Pure and Applied Algebra* (2001) 169-182.
- [3] L. Klingler and A. Omairi, Unique decomposition into ideals for commutative Noetherian rings, *Journal of Pure and Applied Algebra* (2020) 224(9):106364.
- [4] B. Ay Saylam and E. Gurbuz, Unique decomposition into regular ideals for Marot rings.

Rational Groups whose Character Degree Graphs are Disconnected

Gamze Akar

İstinye University
gamze.akar@istinye.edu.tr

Co-author: Temha Erkoç
erkoctemha@gmail.com

Throughout this talk, G is a finite group and $\text{Irr}(G)$ is the set of all irreducible complex characters of G . The set of all irreducible complex character degrees of G is denoted by $\text{cd}(G)$ so that $\text{cd}(G) = \{\chi(1) : \chi \in \text{Irr}(G)\}$. Let $\rho(G)$ be the set of all primes that divide some irreducible character degrees in $\text{cd}(G)$. The character degree graph of G , denoted by $\Delta(G)$, is the graph whose vertex set is $\rho(G)$. Two distinct vertices p, q in $\rho(G)$ are connected by an edge iff there exists at least one degree $a \in \text{cd}(G)$ such that pq divides a . Manz, Willems and Wolf proved in [2] that the character degree graph $\Delta(G)$ of a finite group G has at most three connected components and if G is solvable, then $\Delta(G)$ has at most two connected components. Pálffy showed in [3] that each connected component of the character degree graph of a solvable group must be a complete graph. In [4], Lewis classified all solvable groups having disconnected character degree graphs. Later, Lewis and White classified in [5] nonsolvable groups whose character degree graphs are disconnected. A finite group all of whose complex character values are rational is called a rational group. Motivated by these results, in [1], we classify all rational groups whose character degree graphs are disconnected.

Keywords. Rational Groups, Character degree graphs.

References

- [1] T. Erkoç and G. Akar, Rational Groups whose character degree graphs are disconnected, *Comptes Rendus Mathématique*, preprint (accepted 24th January 2022).
- [2] O. Manz, W. Willems and T. R. Wolf, The diameter of the character degree graph, *J.Reine Angew. Math.* **402**, 181-198, 1989.
- [3] P.P. Pálffy, On the character degree graph of solvable groups, I:three primes, *Period. Math. Hungar*, **36**, 61-65, 1998.
- [4] M.L. Lewis, Solvable groups whose degree graphs have two connected components, *J. Group Theory* **4**, no. 3, 255-275, 2001.
- [5] M.L. Lewis and Donald L. White, Connectedness of degree graphs of nonsolvable groups, *J. Algebra* **266**, no.1, 51-76, 2003.

Mathematical Modeling of Infectious Diseases

Gürçihan Zaman

Dokuz Eylül Üniversitesi

E-mail: `gurcihan.zaman@ogr.deu.edu.tr`

Co-author: Burcu Silindir Yantır

Co-author e-mail: `burcu.silindir@deu.edu.tr`

In this work, we firstly give some basic definitions used in the mathematical models and about the epidemiology of infectious diseases. By making assumptions, we set a model which is called SIR epidemic model, and its mathematical properties is discussed. SIS epidemic model which is a variant of Logistic model is also given. Finally, it is included demography to SIR model, and stability of equilibria of the SIR epidemic model with demography is investigated. This work is based on the book "An introduction to mathematical epidemiology" written by Maia Martcheva. [1]

Keywords. Epidemiology, mathematical models.

References

- [1] MAIA MARTCHEVA, An introduction to mathematical epidemiology, vol. 61 of Text in Applied Mathematics, Springer US, 2015.

Bazzoni's Conjecture and Almost Prüfer Domains

Haleh Hamdi

İzmir Institute of Technology
halehhamdi@iyte.edu.tr

Co-author: Gyu Whan Chang
whan@inu.ac.kr

Bazzoni's conjecture states that if every nonzero locally principal ideal of a Prüfer domain D is invertible, then D is of finite character. This conjecture was proved in [2]. In this talk, we will focus on Bazzoni's conjecture for almost Prüfer domains which states that if every locally finitely generated ideal of an almost Prüfer domain D is finitely generated, then D is of finite character. This result is proved in a more general setting of almost Prüfer v -multiplication domains.

Keywords. Almost Prüfer domain, LPI domain, finite character.

References

- [1] G. W. Chang and H. Hamdi. Bazzoni's conjecture and almost Prüfer domains. *Communications in Algebra*, **47**(7) (2019).
- [2] W. C. Holland, J. Martinez, W. W. McGovern, and M. Tesemma. Bazzoni's conjecture, *Journal of Algebra* **320**(4) (2008).

Mathematical Behavior of Solutions for a Delayed Hyperbolic-Type Equations

Hazal Yüksekaya

University of Dicle
hazally.kaya@gmail.com

Co-author: Erhan Pişkin
episkin@dicle.edu.tr

Controlling the behavior of solutions for partial differential equations with time delay effects has become an active research area. Generally, delay effects occur in many applications and practical problems such as physical, chemical, biological, thermal and economics. In many cases, delay is a source of instability, even an arbitrarily small delay may destabilize a system which is uniformly asymptotically stable in the absence of delay unless additional conditions or control terms have been used. In this work, we study a delayed hyperbolic-type equation. Under appropriate conditions, we prove the mathematical behavior of solutions like existence, blow up, decay.. etc.

Keywords. Delay, Hyperbolic-type equation, Mathematical behavior of solutions.

References

- [1] S. Antontsev, J. Ferreira, E. Pişkin, H. Yüksekaya and M. Shahrouzi, Blow up and asymptotic behavior of solutions for a $p(x)$ -Laplacian equation with delay term and variable exponents, *Electron. J. Differ. Equ.*, **1**-20, 2021.
- [2] M. Kafini and S.A. Messaoudi, Local existence and blow-up of positive-initial-energy solutions of a nonlinear wave equation with delay, *Nonlinear Stud.*, **27**(3), 865-877, 2020.
- [3] S. Nicaise and C. Pignotti, Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks, *SIAM J. Control Optim.*, **45**(5), 1561-1585, 2006.
- [4] E. Pişkin and H. Yüksekaya, Nonexistence of global solutions of a delayed wave equation with variable-exponents, *Miskolc Math. Notes*, **22** (2), 841-859, 2021. 22 (2), 841-859, 2021.
- [5] H. Yüksekaya, E. Pişkin, S. M., Boulaaras and B. B. Cherif, (2021). Existence, Decay, and Blow-Up of Solutions for a Higher-Order Kirchhoff-Type Equation with Delay Term. *Journal of Function Spaces*, 2021.

On 1-Absorbing Intuitionistic Fuzzy Ideals of Commutative Semirings

İlayda Kaplan

Yıldız Technical University
kaplanilayda53@gmail.com

Co-author: E. Mehmet Özkan
mozkan@yildiz.edu.tr

The goal of this research is to look at the algebraic structure of 1-absorbing ideals and how they may be applied to fuzzy sets, as well as the linkages and algebraic features that exist between them. The 1-absorbing intuitionistic fuzzy ideal is examined in this study as an addition to the literature. In this research, intuitionistic fuzzy 1-absorbing ideals are defined, and 1-absorbing ideals and intuitionistic fuzzy sets are used to illustrate instances and theorems.

Keywords. Fuzzy sets, fuzzy ideals, 1-absorbing ideals, 1-absorbing fuzzy ideals, 1-absorbing intuitionistic fuzzy ideals.

References

- [1] L.A.Zadeh, Fuzzy sets, Inf. Control 8 (1965), 338-353.
- [2] A. Badawi and E. Yetkin, On 1-absorbing primary ideals of commutative rings, J. Algebra Appl. (2020), 2050111.
- [3] A.Badawi, On 2-absorbing ideals of commutative rings, Bull. Austral. Math. Soc., 75(2007), 417-429.
- [4] K.Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets Syst. 20 (1986), 87-96.
- [5] K.Hur, H.W.Kang and H.K.Song, Intuitionistic fuzzy subgroups and subrings, Honam Math J. 25 (2003), 19-41.
- [6] M.F.Marashdeh, A.R.Salleh, Intuitionistic fuzzy rings, Int. Journal Algebra 5 (2011), 37-47.

Existence and Stabilization of Solutions of a Linear Viscoelastic Wave Equation with Neumann Boundary Condition

İdem Susuzlu

İzmir Institute of Technology
idemsusuzlu@iyte.edu.tr

Co-author: Türker Özşarı
turker.ozsari@bilkent.edu.tr

We investigate the linear viscoelastic wave equation on a bounded domain $\Omega \subset \mathbb{R}^n$ subject to an inhomogeneous Neumann manipulation activated on a component of the boundary. First, we establish global existence results by separating problem into the two parts. In the first part, we consider usual wave equation with the given initial and boundary data using dynamic extension method. In the second part, we consider viscoelastic part of the equation which also depends on the solution of the first part. We construct approximate solutions by using Faedo-Galerkin method and then use a priori estimates to prove the existence of the weak solutions. Afterwards, we prove uniform stabilization with decay rate determined by behavior of Neumann boundary data and the relaxation function.

Keywords. Viscoelastic wave equations, Relaxation function, Inhomogeneous Neumann boundary data, Faedo-Galerkin method, Approximate solutions, Existence, Stabilization.

References

- [1] Dafermos, C. M., Asymptotic stability in viscoelasticity, Arch. Ration. Mech. Anal. 37, 297-308 (1970).
- [2] David R. Pitts and Mohammad A. Rammaha, Global existence and non-existence theorems for nonlinear wave equations, Indiana Univ. Math. J. 51 (2002), no. 6, 1479-1509.
- [3] Cavalcanti, M. M., Domingos Cavalcanti, V. N. and Soriano, J. A., Exponential decay for the solution of semilinear viscoelastic wave equations with localized damping, Electron. J. Differential Equations 2002(44), 1-14 (2002).
- [4] S.A. Messaoudi, General decay of solutions of a weak viscoelastic equation, Journal of Mathematical Analysis and Applications 36 (2008), 1457-1467.

Stabilization of Higher-Order Schrödinger Equation Using Right Endpoint Controllers

Kemal Cem Yılmaz

İzmir Institute of Technology
cemyilmaz@iyte.edu.tr

Co-author: Türker Özsarı
turker.ozsari@bilkent.edu.tr

Standard backstepping method fails for stabilizing third order evolutionary equations posed on a finite interval, where a single homogeneous boundary condition imposed at one end and control input(s) acting from the other hand. This issue was first addressed in Korteweg de-Vries equation [1] as an open problem. Later it was treated in the case of uncritical domains [2] and in the case of critical domains [3]. The same issue also exists for higher-order Schrödinger equation.

In this talk, we design a feedback controller by modifying the backstepping strategy in such a way that, corresponding closed-loop system is exponentially stable with respect to L^2 -norm both for the case of critical and uncritical lengths of intervals. From the wellposedness point of view, this process requires to prove some regularity estimates for an associated initial-boundary value problem with inhomogeneous boundary data. At the end of the talk, we present a numerical simulation verifying our theoretical result.

Keywords. Higher-order Schrödinger equation, backstepping, boundary controller, observer, stabilization.

References

- [1] E. Cerpa and J.-M. Coron. Rapid stabilization for a Korteweg-de Vries equation from the left Dirichlet boundary condition, *IEEE Trans. Automat. Control* **58**(7) (2013).
- [2] J.-M. Coron and Q. Lü. Local rapid stabilization for a Korteweg-de Vries equation with a Neumann boundary control on the right, *J. Math. Pures Appl.* **102**(6) (2014).
- [3] T. Özsarı and A. Batal. Pseudo-backstepping and its application to the control of Korteweg-de Vries equation from the right endpoint on a finite domain, *SIAM J. Control Optim.* **57**(2) (2019).

3 Noktalı Yol Grafin Karışık Genişlemelerinin Spektral Karakterizasyonu

Kübra Akyıl

Nevşehir Hacı Bektaş Veli Üniversitesi
akyilkubraa@gmail.com

Co-author: Hatice Topcu
haticekamittopcu@gmail.com

Bir G grafinın karışık genişlemesi, G nin herbir noktasının ya bir klik ya da bir ko-klik ile yer değiştirilmesiyle elde edilen H grafidır, öyle ki H grafinda bulunan bu klik ve ko-kliklerdeki noktalar G deki herhangi iki nokta arasındaki komşuluk ilişkisini devam ettirir. Özel olarak G grafi üç noktalı yol graf seçildiğinde, H ın 0 ve -1 e eşit olmayan en fazla üç özdeğeri olur. Bu durum, H grafinın karakterize edilebilmesine imkan sağlayan en temel özelliğini oluşturur [1, 2, 3].

Anahtar Kelimeler. Graf matrisleri, graf spektrumu, karışık genişleme

Kaynakça

- [1] W.H. Haemers. Spectral characterizations of mixed extensions of small graphs, *Discrete Mathematics* **342**(10) (2019).
- [2] H. Topcu, S. Sorgun and W. H. Haemers. The graphs cospectral with the pineapple graph, *Discrete Applied Mathematics* **269**(1) (2019).
- [3] W. H. Haemers, S. Sorgun and H. Topcu. On the spectral characterization of mixed extensions of P_3 , *The Electronic Journal of Combinatorics* **26**(3) (2019).

Metrik Uzaylar Üzerinde Bazı Sabit Çember Teoremleri ve Örnekleri

Kübra Karaağaç

Balıkesir Üniversitesi
kubra.mergen@gmail.com

Co-author: Nihal Taş
nihaltas@balikesir.edu.tr

Sabit noktaların varlığı ve özellikleri ile ilgili teoremler sabit nokta teoremi olarak çalışılmaktadır. Son zamanlarda, sabit nokta kümesine geometrik bir yaklaşım ile sabit çember problemi sabit nokta teorisinin bir genellemesi olarak bırakılmıştır. Biz de bu konuşmada, iki farklı yardımcı fonksiyon yardımıyla sabit çember problemine varlık teoremleri elde ettik. Bu amaç içinde Caristi tipindeki daralma koşulundan yararlanarak, literatürde var olan bazı sonuçları genelledik. Elde ettiğimiz sonuçları gerçekleyen çeşitli örnekler verdik. Ayrıca, sabit çemberlerin tekliği için Khan ve Brianciarı tipinde daralma koşulları ile teklik teoremleri elde ettik.

Keywords. Sabit nokta, sabit çember, varlık teoremi, teklik teoremi.

References

- [1] A. Branciari. A Fixed Point Theorem for Mappings Satisfying a General Contractive Condition of Integral Type, *International Journal of Mathematics and Mathematical Sciences* **29** (2002), Article ID 641824, 6 pages.
- [2] J. Caristi. Fixed Point Theorems for Mappings Satisfying Inwardness Conditions, *Transactions of the American Mathematical Society* **215** (1976), 241-251.
- [3] M. S. Khan, M. Swaleh and S. Sessa. Fixed Point Theorems by Altering Distances Between the Points, *Bulletin of the Australian Mathematical Society* **30** (1984), 1-9.
- [4] N. Y. Özgür and N. Taş. Some Fixed-Circle Theorems on Metric Spaces, *Bulletin of the Malaysian Mathematical Sciences Society* **42**(4) (2019), 433-449.

İşaretli Grafların Spektral Özellikleri

Meliha Akcan

Nevşehir Hacı Bektaş Veli Üniversitesi
melihakcn93@gmail.com

Co-author: Hatice Topcu
haticekamittopcu@gmail.com

Basit bir G grafinın her bir kenarının $+1$ ya da -1 ile işaretlenmesiyle elde edilen H grafinı işaretli graf denir, bu tanım ilk olarak [4] te Zaslavsky tarafından verilmiştir. Son zamanlarda, işaretli grafların spektrumları, graf spektrası çalışan araştırmacılar tarafından çok fazla ilgi görmektedir[1, 2, 3]. Bu durumun temel motivasyonu, işaretli grafların spektral teorisinin, işaretli grafların spektral teorisinin zarif bir şekilde genelleştirilmesi biçiminde olmasıdır. Öte yandan, işaretli graflar, asıl olarak dengeli işaretli grafların özel bir durumunu teşkil etmektedir.

Anahtar Kelimeler. Graf matrisleri, graf spektrumu, işaretli graf

Kaynakça

- [1] T. Zaslavsky. Signed graphs, *Discrete and Applied Mathematics* **1**(4) (1982).
- [2] F. Belardo, S.M. Cioaba, J. Koolen and J. Wang. Open problems in the spectral theory of signed graphs, *The Art of Discrete and Applied Mathematics* **1**(2) (2018).
- [3] T. Zaslavsky. Negative and positive circles in signed graphs: a problem collection, *AKCE International Journal of Graphs and Combinatorics* **15**(1) (2018).
- [4] W. H. Haemers and H. Topcu. On signed graphs with at most two eigenvalues unequal to ∓ 1 , <https://doi.org/10.48550/arXiv.2109.02522> (2021).

Sıra Norm Sürekli Operatörlerin Latis Yapısı Üzerine

Merve Bülbül

Gazi Üniversitesi
merve4214@gmail.com

Co-author: Birol Altın
birola@gazi.edu.tr

Kazem Hanghnejad Azar, Ali Rıza Jalili ve Mohammad Farsbaf Moghili isimli yazarlar Positivity dergisinde 2021 yılında ‘order-to-topology continuous operators’ isimli makalelerinde sıra-topolojik-sürekli operatörü tanıtmışlardır. Yazarlar bu çalışmasında E bir Riesz uzayı ve F bir normlu Riesz uzayı olmak üzere E den F ye tanımlı her sıra norm sürekli operatörünün modülünün olup olmadığını açık problem olarak bırakmışlardır. Biz bu çalışmada bu problemin genelde doğru olmadığını gösteren bir örnek inşa edeceğiz ayrıca yukardaki açık probleme bazı özel şartlar yükleyerek T nin modülünün var ve sıra norm sürekli olduğu sonucu verilecektir.

Roth's Theorem on Arithmetic Progressions

Mustafa Kutay Kutlu

Dokuz Eylül University
mustafakutay.kutlu@ogr.deu.edu.tr

Co-author: Assoc. Prof. Dr. Engin Mermut
engin.mermut@deu.edu.tr

An *arithmetic progression of length k* (k -AP) is a sequence of k -numbers such that each difference between two consecutive terms is the same constant. Finding long arithmetic progressions in certain subsets of integers is at the center of mathematics in the last century. In this talk, we will first give a quick survey of remarkable results on this topic. In particular, we will focus on Roth's theorem on arithmetic progressions. Finally, we will also talk about recent significant results.

Keywords. Arithmetic progression, Roth's theorem.

References

- [1] Endre Szemerédi. On sets of integers containing no elements in arithmetic progression, *Acta Arithmetica* **27**(199-245).
- [2] Klaus Roth. On certain sets of integers, *J. London Math Soc.* **28**(104-109).
- [3] Van der Waerden. Beweis Archief voor Wiskunde, *Nieuw Arch. Wiskd.* **15**(212-216).

New Infinite Rank Summands of Homology Cobordism Group

Oğuz Şavk

Boğaziçi University, İstanbul and Max-Planck-Institut für Mathematik, Bonn
 oguz.savk@boun.edu.tr, savk@mpim-bonn.mpg.de

Co-author: Çağrı Karakurt
 cagri.karakurt@boun.edu.tr

Since the 1980s, the homology cobordism group $\Theta_{\mathbb{Z}}^3$ has been a central object in low-dimensional topology. Although its algebraic structure is still mysterious, the recent achievement of Dai, Hom, Stoffregen, and Truong indicates that it is very complicated. They prove that $\Theta_{\mathbb{Z}}^3$ has an infinite rank summand \mathbb{Z}^{∞} generated by the Brieskorn homology spheres $\{X_n = \Sigma(2n+1, 4n+1, 4n+3)\}_{n=1}^{\infty}$, see [1].

We generalize this link of singularity by assuming that $pq + pr - qr = 1$ where p, q and r are pairwise coprime ordered positive integers. Clearly, the parameters of the family $\{X_n\}_{n=1}^{\infty}$ satisfy the latter identity. Then Brieskorn homology spheres $\Sigma(p, q, r)$ can be realized as the boundaries of *almost simple linear graphs*.

We compute Ozsváth-Szabó d -invariants and connected Heegaard Floer homologies of two families of Brieskorn homology spheres $\{Y_n = \Sigma(2n+1, 3n+2, 6n+1)\}_{n=1}^{\infty}$ and $\{Z_n = \Sigma(2n+1, 3n+1, 6n+5)\}_{n=1}^{\infty}$, see [4], [2], and [3] respectively. They are also two specific boundaries of almost simple linear graphs. Using Floer theoretic invariants of Dai, Hom, Stoffregen, and Truong, we show that $\{Y_n\}_{n=1}^{\infty}$ and $\{Z_n\}_{n=1}^{\infty}$ also generate infinite rank summands in the homology cobordism group. We will also compare their homology cobordism classes and we will see that they are not cobordant to each other with an exception $X_1 = \Sigma(3, 5, 7) = Y_1$.

In this talk, our aim is to present the framework of our effective computation procedure and discuss its combinatorial nature with several illustrations. This is joint work with Çağrı Karakurt.

Keywords. Brieskorn spheres, homology cobordism group, Heegaard Floer homology.

References

- [1] Dai, I., Hom, J., Stoffregen, M., Truong, L. (2018). An infinite-rank summand of the homology cobordism group. arXiv preprint arXiv:1810.06145.
- [2] Hendricks, K., Hom, J., Lidman, T. (2021). Applications of involutive Heegaard Floer homology. *Journal of the Institute of Mathematics of Jussieu*, 20(1), 187-224.
- [3] Karakurt, Ç., Şavk, O. (2020). Ozsváth-Szabó d -invariants of almost simple linear graphs. *Journal of Knot Theory and Its Ramifications*, 29(5), 2050029.
- [4] Karakurt, Ç., Şavk, O. (2022). Almost simple linear graphs, homology cobordism and connected Heegaard Floer homology. arXiv preprint arXiv:2204.06597.

Truncation on ℓ -groups

Osman Hamza

Yıldız Technical University
hamzaosman61@gmail.com

In this study, we give the definition of truncation and its properties on ℓ -groups with truncation. Here, the ℓ -group means a divisible abelian lattice-ordered group. Finally, we indicate several operations on the truncated ℓ -groups.

Keywords. truncation, ℓ -group, lattice ordered group

References

- [1] Boulabiar, K., El Adeb, C.: Unitization of Ball truncated -groups. Algebra Univ. 78, 93-104 (2017).
- [2] Boulabiar, K., Mahfoudhi, M.: Unitization of a lattice ordered ring with a truncation, Quaestiones Mathematicae, 43:7, 841-856 (2020).

Metrik Uzaylarda Sabit Çemberler

Özlem Moralı Kızanlık

Balıkesir Üniversitesi
ozlemmorali@outlook.com

Co-author: Nihal Taş
nihaltas@balikesir.edu.tr

Sabit nokta teorisi Stefan Banach zamanından beri bilimin pek çok dalında hem teorik hem de uygulama anlamında kullanılmaktadır. Bu konuşmada, sabit nokta teorisine geometrik bir yaklaşım olarak, metrik uzaylar üzerinde bazı dönüşümlerin sabit çemberler için varlık ve teklik teoremleri vereceğiz. Farklı metrik fonksiyonları kullanarak bir çemberin eleman sayısının değişebileceğini vurgulayıp, alışılmış metrik uzay üzerinde de teoremleri sağlayan örnekler vereceğiz.

Keywords. Sabit nokta, sabit çember, varlık teoremi, teklik teoremi.

References

- [1] S. Banach. Sur Les Oprations Dans Les Ensembles Abstraits et Leur Applications Aux Equations Integrales, *Fundemental Mathematics* **3** (1922), 133-181.
- [2] U. Çelik and N. Özgür. On the Fixed Circle Problem, *Facta Universitatis. Series: Mathematics and Informatics* **35**(5) (2020), 1273-1290.
- [3] L. B. Ćirić. A Generalization of Banach's Contraction Principle, *Proceedings of the American Mathematical Society* **45**(2) (1974), 267-273.
- [4] N. Y. Özgür and N. Taş. Some Fixed-Circle Theorems on Metric Spaces, *Bulletin of the Malaysian Mathematical Sciences Society* **42**(4) (2019), 433-1449.

ωe^* -Ayrırma Aksiyomları

Pınar Şaşmaz

Muğla Sıtkı Koçman Üniversitesi
pinarsasmazz@gmail.com

Co-author: Murad Özkoç
murad.ozkoc@gmail.com

Topolojiler incelidikçe farklı nokta ve farklı kümeleri, farklı komşuluklarla ayırma imkanı doğar. Bu noktada önemli olan ise araştırmacıların küme üzerinde yapmak istediği işe uygun bir topolojik yapı kurmasıdır. Bu sebeple topolojik yapıları bazı ortak özellikleri olan sınıflara ayırmak daha yararlı olmaktadır. Bu noktada ayırma aksiyomlarının önemli bir rolü vardır.

Son yıllarda ayırma aksiyomlarının bazı özel ve genel formları, birçok araştırmacı tarafından tanımlanmış ve çalışılmıştır. Ayrıca açık kümelerin özel ve genel bazı formları, bu çalışmalara ayrı bir önem kazandırmıştır. 1983 yılında M.E. Abd El-Monsef, S.N. El-Deeb ve R.A. Mahmoud tarafından verilen β -açık küme [3] ve 2003 yılında K.Y. Al-Zoubi ve B. Al-Nashef tarafından verilen ω -açık küme [2] kavramlarından yola çıkarak H.H. Aljarrah, M.S.M. Noorani ve T. Noiri tarafından tanımlanan $\omega\beta$ -açık kümeler [1] ile 2015 yılında H.H. Aljarrah, M.S.M. Noorani ve T. Noiri tarafından tanımlanan $\omega\beta$ -ayırma aksiyomları [5] bunun önemli örneklerinden biri olarak karşımıza çıkmaktadır.

Bu çalışmada ise $\omega\beta$ -açık küme kavramından daha genel bir kavram olan ωe^* -açık küme [4] kavramından hareketle $\omega\beta$ -ayırma aksiyomlarından daha zayıf bir kavram olarak karşımıza çıkan ωe^* -ayırma aksiyomları çalışılmıştır. ωe^*-T_0 uzay, $\omega e^*-T_{\frac{1}{2}}$ uzay, ωe^*-T_1 uzay ve ωe^*-T_2 uzay kavramları tanıtılmış ve bu kavramların bazı karakterizasyonları elde edilmiştir. Ayrıca ωe^*-R_0 uzay ve ωe^*-R_1 uzay kavramları tanımlanarak aralarındaki bazı ilişkiler incelenmiştir.

Anahtar Kelimeler. ωe^* -açık küme, ωe^*-T_1 uzay, $\omega e^*-T_{\frac{1}{2}}$ uzay, ωe^*-T_1 uzay, ωe^*-T_2 uzay, ωe^*-R_0 uzay, ωe^*-R_1 uzay

Kaynaklar

- [1] M.E. Abd El-Monsef, S.N. El-Deeb and R.A. Mahmoud, β -open sets and β -continuous mapping, *Bull. Fac. Sci. Assiut Univ.*, **12** (1983), 77-90.
- [2] H.H. Aljarrah, M.S.M. Noorani and T. Noiri, On $\omega\beta$ -open sets, *submitted*.
- [3] H.H. Aljarrah, M.S.M. Noorani and T. Noiri, $\omega\beta$ -separation axioms, *Vasile Alecsandri University of Bacău Faculty of Sciences Scientific Studies and Research Series Mathematics and Informatics*, **25**(1), (2015), 87-106.
- [4] K.Y. Al-Zoubi and B. Al-Nashef, The Topology of ω -open subsets, *Al-Manarah Journal*, **9** (2003), 169-179.

- [5] M. Özkoç and P. Şaşmaz, On contra ωe^* -continuous functions, *Poincare J. Anal. Appl.*, **8**(1(I)), (2021), 51-65.

On 2-Absorbing Vague Fuzzy Ideals of Commutative Semirings

Rabia Ayan

Yıldız Technical University
rabiaayan408@gmail.com

Co-author: E. Mehmet Özkan
mozkan@yildiz.edu.tr

The purpose of this study is to investigate the algebraic structure of 2-absorbing ideals, as well as how they may be applied to fuzzy sets and the connections and algebraic properties that exist between them. This study adds to the literature by looking at the 2-absorbing vague fuzzy ideal. In this study, 2-absorbing vague fuzzy ideals are established, and cases and theorems are shown using 2-absorbing ideals and vague fuzzy sets.

Keywords. Fuzzy sets, fuzzy ideals, 2-absorbing ideals, 2-absorbing fuzzy ideals, 2-absorbing vague fuzzy ideals.

References

- [1] L.A.Zadeh, Fuzzy sets, Inf. Control 8 (1965), 338-353.
- [2] A.Badawi, On 2-absorbing ideals of commutative rings, Bull. Austral. Math. Soc., 75 (2007), 417-429.
- [3] Doebner, H. and Henning, J., eds., Quantum Groups, Springer, Berlin, (1990).
- [4] K.Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets Syst. 20 (1986), 87-96.
- [5] Faddeev, L.D., Les Houches Lectures 1982 - 1984. Elsevier, Amsterdam (1984).
- [6] Jimbo, M., A q-difference analogue of $U(g)$ and the Yang-Baxter equation, Lett. Math. Phys.10, (1981), 63-69.

Dirichlet Boundary Value Problem for the Laplace Equation in the Semi-Infinite Domain via a Boundary Integral Equation Method

Sabahat Defne Plattürk

İzmir Institute of Technology
sabahatplatturk@iyte.edu.tr

Co-author: Olha Ivanyshyn Yaman
olhaivanyshyn@iyte.edu.tr

We propose a method to solve the boundary value problem for the Laplace equation in a doubly connected semi-infinite domain. With the aid of a conformal map the semi-infinite domain is transformed to a doubly connected bounded domain. Then the problem is reduced to a system of linear boundary integral equations by representing the solution to the boundary value problem as a combination of double- and single-layer potentials with a modification that ensures the unique solvability of the system of Fredholm integral equations of the second kind. The system is solved numerically via the Nyström method[1].

Keywords. Boundary value problem, laplace equation, boundary integral equations, nyström method, conformal mapping

References

- [1] Kress, Rainer, Linear integral equations. Vol. 82. Berlin: Springer, 1989.

A Generalization of Injective Modules

Sinem Benli Göral

İzmir Institute of Technology
sinembenli@iyte.edu.tr

Co-authors:

Yusuf Alagöz¹, Engin Büyükaşık²

¹yusuf.alagoz@siirt.edu.tr, ²enginbuyukasik@iyte.edu.tr

Harada [4] introduced the notions of mininjective modules and rings as follows: Let M and N be right R -modules. M is said to be *min- N -injective* if for every simple submodule K of N , and every homomorphism $f : K \rightarrow M$, there exists a homomorphism $h : N \rightarrow M$ such that $h|_K = f$. If we take the right R -module R_R for N , then M is called *mininjective*. On the other hand, according to Harada [5], M is said to be *simple- N -injective* if for every submodule K of N , and every homomorphism $f : K \rightarrow M$ with $f(K)$ simple extends to N . If $N = R_R$, M is called *simple-injective*. The concepts of mininjectivity and simple-injectivity of rings and modules aroused interest and many results on them and their generalizations have been appeared in the literature, see for example [2, 3, 6, 7]. In this talk, we aim to present some new results about the aforementioned modules that are recently appeared in [1].

Keywords. Quasi-Frobenius rings, injective, mininjective, simple-injective.

References

- [1] Y. Alagöz, S. Benli-Göral and E. Büyükaşık, On simple-injective modules, *Jour. Alg. Appl.* (accepted).
- [2] I. Amin, Y. Fathi and M. F. Yousif, Strongly simple-injective rings and modules, *Alg. Coll.* **15(1)** (2008) 135–144.
- [3] I. Amin, M. F. Yousif and N. Zeyada, Soc-injective rings and modules, *Comm. Algebra* **33** (2005) 4229–4250.
- [4] M. Harada, Self mini-injective rings, *Osaka J. Math.* **19(2)** (1982) 587–597.
- [5] M. Harada, Note on almost relative projectives and almost relative injectives, *Osaka J. Math.* **29** (1992) 435–446.
- [6] L. Mao, Rings related to mininjective and min-flat modules, *Comm. Algebra* **37** (2009) 3586–3600.
- [7] M. F. Yousif and Y. Zhou, *FP*-injective, simple-injective, and quasi-Frobenius rings, *Comm. Algebra* **32** (2004) 2273–2285.

Classification of Orlicz Spaces equipped with s-norms

Şeyma Yaşar

Gebze Technical University
seymayasar@gtu.edu.tr

Co-author: Serap Öztıp, Esra Başar, Badik Hüseyin Uysal
oztops@istanbul.edu.tr esra.basar@yeditepe.edu.tr
huseyinuyisal@istanbul.edu.tr

Orlicz Spaces equipped with s norms introduced by M. Wisła. L_s^Φ is Orlicz spaces equipped with s-norms. In this study, our aim is to classify s norms with respect to σ_s numbers which is defined by $\sigma_s := \sup\{u \geq 0 : s(u) = 1\}$. This classification helps us for investigate the s norms more easily.

Keywords. Orlicz space, Extreme points, Compact operators

References

- [1] Chen, S. and Wisła, M., Extreme compact operators from Orlicz spaces to $C(\Omega)$, *Comment. Math. Univ. Carolin.*, 34(1), (1993)
- [2] Dunford, N. and Schwartz, J.T., *Linear Operators I: General Theory*, Interscience, (1958).
- [3] M. Wisła, Orlicz spaces equipped with s-norms, *Journal of Mathematical Analysis and Applications*, **483**(2), 2020
- [4] Rao, M.M. and Ren, Z.D., *Theory of Orlicz Spaces*, Marcel Dekker,(1991).

Spin Structures on Real Bott Manifolds

Şule Kılıçaslan

Dokuz Eylül University
sule.kilicaslann@gmail.com

Co-author: Assoc. Prof. Dr. Ash Güçlükan İlhan
asli.ilhan@deu.edu.tr

A real Bott tower of height n is a sequence of real projective bundles

$$B_n \longrightarrow B_{n-1} \longrightarrow \cdots \longrightarrow B_1 \longrightarrow \{point\} \quad (2)$$

such that for each $j \in \{1, 2, \dots, n\}$, $B_j \rightarrow B_{j-1}$ is the projectivization of the Whitney sum of a real line bundle L_j and the trivial line bundle over B_{j-1} [1]. The manifold B_n is called a real Bott manifold.

In this talk, we first give necessary and sufficient conditions for the existence of a Spin structure on real Bott manifolds. This result is due to [2]. Then we discuss some results on the number of real Bott manifolds with Spin structures for particular n 's that we obtained using SageMath.

Keywords. Acyclic Digraphs, Real Bott Manifold

References

- [1] M. Grossberg and Y. Karshon. Bott towers, complete integrability, and the extended character of representations, *Duke Math. J.* 76 23-58 (1994).
- [2] R Dsouza. On the topology of real Bott manifolds, *Indian Journal of Pure and Applied Mathematics* 9 (4) 743-763 (2018).

Asymptotic Behavior of Solutions for p-biharmonic Parabolic Equation with Logarithmic Nonlinearity

Tuğrul Cömert

Dicle University
tugrulcomertt@gmail.com

Co-author: Erhan Pişkin
episkin@dicle.edu.tr

It is possible that many of the problems we encounter in daily life can be modeled with ordinary or partial differential equations. These problems can be related to many fields from physics to chemistry, from biology to medical science. However, it is often not possible to find an exact solution to every modeled problem. In order to find an approximate solution for these problems or at least to have an idea about the behavior of the solution, it has become necessary to limit the equation to some conditions. For this reason, it is aimed to find well-defined solutions by adding initial and boundary conditions to the problems. In order to find a well-defined solution, the first step is to investigate the existence of the solution, the uniqueness of the solution, if any, and its dependence on the initial data. The existence of the solution and the asymptotic behavior of the equation that has proven to be unique, that is, the behavior of the solution in a finite or infinite time, it allows us to have an idea even if the solution is not known exactly. This paper investigate the p-biharmonic parabolic equation with logarithmic nonlinearity in a bounded domain. Firstly, we consider the global existence of solutions. Later, we investigate the decay of solutions.

Keywords. p-biharmonic parabolic equation, Asymptotic behavior, Logarithmic nonlinearity.

References

- [1] A. J. Hao and J. Zhou, Blow up, extinction and non-extinction for a nonlocal p-biharmonic parabolic equation, *Applied Mathematics Letters* **64** (2017), 198-204.
- [2] C. Liu and J. Guo, Weak solutions for a fourth order degenerate parabolic equation, *Bulletin of the Polish Academy of Sciences, Mathematics* **54**(1) (2006), 27-39.
- [3] L. C. Nhan and L. X. Truong, Global solution and blow-up for a class of pseudo p-Laplacian evolution equations with logarithmic nonlinearity, *Computers and Mathematics with Applications* **73** (2017), 2076-2091.
- [4] J. Wang and C. Liu, p-biharmonic parabolic equations with logarithmic nonlinearity, *Electronic Journal of Differential Equations* **2019**(8) (2019), 1-18.

On the Depth and Reflexivity of Modules

Uyen Enni Le

West Virginia University
hle1@mix.wvu.edu

In 1994, Huneke and Wiegand introduced the Depth Formula, but in 2007, they announced in an erratum that one of the conclusions in the depth formula theorem is flawed due to an incorrect convention for the depth of the zero module. Not until 2019 was the claim proven to be false by Celikbas and Takahashi. In this talk, we determine new conditions under which the depth formula theorem holds true, i.e., if the tensor products of finitely generated modules over hypersurface rings is reflexive, then both of their factors are reflexive. Finally, we will discuss some examples to show why these results are new and necessary.

Categorical Methods in the Representation Theory of Artin Algebras

Victor Blasco

Dokuz Eylül University
victorblasco98@gmail.com

In this talk we will introduce the basics of the representation theory of Artin Algebras. In particular, the notion of right minimal morphisms will help us to prove the existence of a projective cover for any finitely generated module. We will also characterize the indecomposable projective modules over these algebras.

Keywords. Artin algebras, representation theory

References

- [1] M. Auslander, I. Reiten, S.O. Smalø. Representation Theory of Artin Algebras, *Cambridge University Press*, 1995

Applications of Cayley Digraphs to Waring's Problem and Sum-Product Formulas

Yeşim Demiroğlu Karabulut

California Institute of Technology
yesim@caltech.edu

In this talk, we first present some elementary new proofs (using Cayley digraphs and spectral graph theory) for Waring's problem over finite fields, and explain how in the process of re-proving these results, we obtain an original result that provides an analogue of Sárközy's theorem in the finite field setting (showing that any subset E of a finite field \mathbb{F}_q for which $|E| > \frac{qk}{\sqrt{q-1}}$ must contain at least two distinct elements whose difference is a k^{th} power). Once we have our results for finite fields, we can apply some classical mathematics to extend our Waring's problem results to the context of general (not necessarily commutative) finite rings. In the second half of our talk, we talk about some sum-product formulas related to matrix rings over finite fields, which can again be proven using Cayley digraphs and spectral graph theory in an efficient way.

Keywords. Waring's problem, Cayley digraphs, finite fields.

Döngüsel Çizgelerin Ricci Eğrilikleri

Yonca Ünver

Amasya Üniversitesi
yonca-gs@hotmail.com

Co-author: Mehmet Dağlı
mehmet.dagli@amasya.edu.tr

Lin, Lu ve Yau tarafından tanıtılan (kaba) Ricci eğriligi, lokal olarak sonlu çizgeler üzerinde bir izomorfizm invariantı sunmaktadır. Bu nedenle, çizge ailelerinin Ricci eğriliklerine ait bir katalog oluşturmak önemlidir. Eğriliklerin hesaplanmasında kullanılan önemli araçlardan biri Eşleşme Koşuludur. Bu çalışmanın amacı, eşleşme koşulunu kullanarak Ricci eğriligi sabit olan döngüsel çizge aileleri oluşturmaktır. Burada kullanılan yöntem, Cayley çizgeleri için de kullanılabilir.

Keywords. Çizge, döngüsel çizge, Ricci eğriligi, eşleşme koşulu.

References

- [1] B. B. Bhattacharya ve S. Mukherjee. Exact and asymptotic results on coarse Ricci curvature of graphs, *Discrete Math.* **338**(1) (2015).
- [2] J. D. H. Smith. Ricci curvature, circulants, and a matching condition, *Discrete Math.* **329** (2014).
- [3] M. Dağlı, O. Ölmez ve J.D.H. Smith. Ricci curvature, circulants, and extended matching conditions, *Bull. Korean Math. Soc.* **56**(1) (2019).
- [4] Y. Lin, L. Lu ve S.-T. Yau. Ricci curvature of graphs, *Tohoku Math. J.* **63**(4) (2011).
- [5] Y. Ollivier. Ricci curvature of Markov chains on metric spaces, *J. Funct. Anal.* **256**(3) (2009).

Time-Evolution of Cauchy-Euler Type Quantum Oscillator in the Presence of Variable Magnetic and Electric Fields

Zehra Çayıç

İzmir Institute of Technology
zehracayic@iyte.edu.tr

Co-author: Şirin Atılğan Büyükaşık
sirinatilgan@iyte.edu.tr

We introduce an exactly solvable Cauchy-Euler type quantum parametric oscillator in the presence of time-variable external fields. The corresponding evolution problem is solved by using Wei-Norman Lie algebraic approach [1], [3]. Then, time-evolution of the eigenstates and coherent states [2], is found explicitly in terms of solutions to the corresponding system of coupled classical equations of motion. Also, squeezing properties of the wave packets [4], and their trajectories in the two-dimensional configuration space are discussed according to the influence of the time-variable parameters and external fields.

References

- [1] J. Wei, E. Norman. Lie algebraic solution of linear differential equations, *J.Math.Phys.* **4**, 575 (1963).
- [2] Roy J. Glauber. Coherent and incoherent states of the radiation field, *Phys. Review* **131**, 2766(1963).
- [3] Ş. A. Büyükaşık and Z. Çayıç. Exactly solvable Hermite, Laguerre, and Jacobi type quantum parametric oscillators, *J. Math. Phys.* **57**, 122107 (2016).
- [4] Ş. A. Büyükaşık and Z. Çayıç. Time-evolution of squeezed coherent states of a generalized quantum parametric oscillator, *J. Math. Phys.* **60**, 062104 (2019).

İki Değişkenli q -Bleimann, Butzer ve Hahn Operatörlerinin Yaklaşım Oranları

Zeynep Kara

Gazi Üniversitesi
zeynepkara1930@gmail.com

Co-author: Prof.Dr.Ogün Doğru
ogun.dogru@gazi.edu.tr

Bu çalışmada Lineer pozitif operatör tanımından başlanarak süreklilik modülü, Lipschitz sınıfı fonksiyonlar ve operatör dizileri için düzgün yakınsaklık kavramı ve bununla birlikte Korovkin Teoremi ele alınmıştır. $C_B[0, \infty)$ uzayının bir alt uzayı olan H_w uzayı tanımlanmış ve Bleimann, Butzer ve Hahn operatör dizisinin H_w uzayındaki fonksiyonlar için $[0, \infty)$ aralığında düzgün yakınsaklığı incelenmiştir. Daha sonra bu operatörün n ye göre monotonluğu bölünmüş farklar yardımıyla verilmiştir. q tamsayısına dayalı Bleimann, Butzer ve Hahn operatörlerinin tanımı [1] verilip bu operatör dizisinin düzgün yakınsaklığı incelenmiştir. Ayrıca bu operatör dizisinin yaklaşım hızı süreklilik modülü ve Lipschitz tipli maksimal fonksiyonlar yardımıyla değerlendirilmiştir. Ardından q tamsayısına dayalı Bleimann, Butzer ve Hahn operatörlerinin iki değişkenli hali tanımı [2] verilerek bu operatörlerin reel uzayın sınırlı ve sürekli bir alt uzayında sürekli bir fonksiyona düzgün yakınsadığı gösterilmiştir. Ayrıca bu operatör dizisinin yaklaşım hızı süreklilik modülü ve Lipschitz tipli maksimal fonksiyonlar yardımıyla değerlendirilmiştir. Daha sonra özgün olarak q - Bleimann, Butzer ve Hahn operatörlerinin Voronovskaya tipli asimptotik yaklaşımı ve iki değişkenli q - Bleimann, Butzer ve Hahn operatörlerinin Voronovskaya tipli asimptotik yaklaşımı elde edilmiştir.[2]

Anahtar Kelimeler. Bleimann Butzer Hahn operatörü , lineer pozitif operatörler,Korovkin teoremi, q -analiz, süreklilik modülü, lipschitz sınıfı

Kaynakça

- [1] Aral, A. and Doğru, O. Bleimann Butzer and Hahn operators based on q -integers. *Journal of Inequalities and Applications*, Art. ID 79410; 1-12. (2007).
- [2] Altin, A., Doğru, O., On the Approximation Properties of Bivariate Bleimann, Butzer and Hahn Operators. *WSEAS Trans. Math.* 4(4), 327-332. (2005)

LIST OF PARTICIPANTS

Abdulkaki Aşur	Dokuz Eylül University
Ahmet Çolak	İzmir Institute of Technology
Ali Öksüz	İzmir Institute of Technology
Alican Yaşar	İzmir Institute of Technology
Aliye Yiğit	Hacettepe University
Arjin Budak	İzmir Institute of Technology
Asgar Jamneshan	Koç University
Aslı Güçlükan İlhan	Dokuz Eylül University
Aygül Koçak Özvarol	İzmir Institute of Technology
Aykut Alkın	İzmir Institute of Technology
Aylin Bozacı Serdal	İzmir Institute of Technology
Ayten Erilmez	İzmir Institute of Technology
Ayça İleri	Dokuz Eylül University
Ayşe Aybala Doğan	İzmir Institute of Technology
Ayşe Fidan	Dicle University
Ayşegül Çetin	İzmir Institute of Technology
Azem Berivan Adıbelli	İzmir Institute of Technology
Badik Hüseyin Uysal	İstanbul University
Banu Pazar Varol	Kocaeli University
Barış Çiçek	İzmir Institute of Technology
Başak Ay Saylam	İzmir Institute of Technology
Berrak Özgür	İzmir Democracy University
Betül Kızılkaya	İzmir Institute of Technology
Betül Tolgay	Middle East Technical University
Burak Erdoğan	University of Illinois at Urbana-Champaign
Burak Gürel	Boğaziçi University
Burak Turfan	İzmir Democracy University
Burcu Silindir	Dokuz Eylül University
Buğurcan Dişibüyük	Dokuz Eylül University
Canan Özeren	Dokuz Eylül University
Celal Cem Sarıoğlu	–
Cihan Okay	Bilkent University
Cihan Sahillioğulları	İzmir Institute of Technology
Cihat Aydın	Yıldız Teknik University
Çağatay Altuntaş	İstanbul Technical University
Çetin Dişibüyük	Dokuz Eylül University
Çınar Eren Arslan	Koç University
Derya Özdemir	İzmir Institute of Technology
Dilan Cantosun	İzmir Institute of Technology
Doğa Can Sertbaş	Çukurova University
Duygu Uysal	İzmir Institute of Technology
Ebru Nisa Koç	Çukurova University
Ece Hazal Korkmaz	İstanbul Technical University
Ecem Tıver	Dokuz Eylül University
Eda Nur Uzun	İzmir Institute of Technology
Ege Tamcı	İzmir Institute of Technology
Elif Ergül	İzmir Institute of Technology
Elif Güner	Kocaeli University
Elifnur Demirkol	İzmir Institute of Technology
Emre Akharman	İzmir Institute of Technology
Emre Yılmaz	Julius-Maximilians-Universität Würzburg
Engin Başakoğlu	Boğaziçi University
Engin Büyükaşık	İzmir Institute of Technology
Engin Mermut	Dokuz Eylül University
Engin Özkan	Ted University
Erdem Şafak Öztürk	Ankara University
Eren Şen	Middle East Technical University
Eren Canan	İzmir Institute of Technology
Ergün Yalçın	Bilkent University
Esmâ Dirican Erdal	Yeditepe University
Esmeray Yevimli	İzmir Institute of Technology
Esra Altıntaş	Ege University
Esra Başar	Yeditepe University

Ezgi Gürbüz	İzmir Institute of Technology
Ezgi Tutar	İzmir Institute of Technology
Faruk Temur	İzmir Institute of Technology
Fatih Çelik	Vefa Lisesi
Fatih Erman	İzmir Institute of Technology
Fatih İdiz	İzmir Institute of Technology
Fatma Zürcacı Yetiş	İstanbul Technical University
Fikriye Nuray Yılmaz	Gazi University
Gamze Alkaya	Gazi University
Gamze Akar	İstinye University
Gökhan Karahanoğulları	İzmir Institute of Technology
Gökhan Şahan	İzmir Institute of Technology
Gülter Budakçı	Dokuz Eylül University
Gürcihan Zaman	Dokuz Eylül University
Hakan Şanal	Dokuz Eylül University
Haleh Hamdi	İzmir Institute of Technology
Halil Oruç	Dokuz Eylül University
Hatice Topcu	Nevşehir Hacı Bektaş Veli University
Haydar Baran Yurtsever	İzmir Institute of Technology
Haydar Göral	İzmir Institute of Technology
Hazal Yüksekaya	Dicle University
Hikmet Burak Özcan	İzmir Institute of Technology
İdem Susuzlu	İzmir Institute of Technology
İlayda Kaplan	Yıldız Teknik University
İrem Baydoğan	Middle East Technical University
İrem Zengin	İzmir Democracy University
İzzet Coşkun	University of Illinois at Chicago
Jan Stovicek	Charles University
Joshua Zahl	University of British Columbia
Julian Weigt	Aalto University
Kemal Cem Yılmaz	İzmir Institute of Technology
Kübra Akyl	Nevşehir Hacı Bektaş Veli University
Kübra Karaağaç	Balıkesir University
Leyla Akbalık	İzmir Institute of Technology
Mehmet Kaymak	İzmir Institute of Technology
Melda Duman	Dokuz Eylül University
Melek Sofyaloğlu	Ankara Hacı Bayram Veli University
Meliha Akcan	Nevşehir Hacı Bektaş Veli University
Meltem Adıyaman	Dokuz Eylül University
Meltem Uzun	İstanbul Gelişim University
Meltem Güllüsaç	Dokuz Eylül University
Mert Akman	İzmir Institute of Technology
Merve Bülbül	Gazi University
Merve Kahraman Arıman	Dokuz Eylül University
Musa Elsıkma	İzmir Institute of Technology
Mustafa Kırçalı	Middle East Technical University
Mustafa Kutay Kutlu	Dokuz Eylül University
Mustafa Özel	Dokuz Eylül University
Müge Diril	İzmir Institute of Technology
Nazlıcan Ata	İzmir Institute of Technology
Negar Bakhshi	Farhangyan University
Neslihan Güneş	İzmir Institute of Technology
Nesrin Yakar	İzmir Institute of Technology
Nihal Taş	Balıkesir University
Nihan Tanısalı	Koç University
Nurdan Poyraz	–
Oktay Pashaev	İzmir Institute of Technology
Olha Ivanyshyn Yaman	İzmir Institute of Technology
Onur Aytay	İzmir Institute of Technology
Orhun Kara	İzmir Institute of Technology
Osman Hamza	Yıldız Teknik University
Oğuz Şavk	Boğaziçi University & Max Planck Institute for Mathematics
Oğuz Yılmaz	İzmir Institute of Technology
Özgür Ege	Ege University
Özlem Irmak Demir	İzmir Institute of Technology
Özlem Moralı Kızanlık	Balıkesir University

Pınar Eroğlu
 Pınar Şaşmaz
 Rabia Atasoy
 Rabia Ayan
 Refail Alizade
 Sabahat Defne Plattürk
 Sabri Kaan Gürbüz
 Sadık Eyidoğan
 Salahattin Özdemir
 Saliha Kıvanç
 Seda Arslan
 Selçuk Demir
 Selin Erdal
 Sena Nur Karataş
 Seven Burçin Çellek
 Seçil Gergün
 Shodijon Ismoilov
 Sinem Benli Göröl
 Sinem Onaran
 Sueda Nur Uysal
 Süleyman Cengizci
 Sultan Eylem Toksoy
 Şermin Çam Çelik
 Şeyhmus Can
 Şeyma Yaşar
 Şirin Atılğan Büyükaşık
 Şule Kılıçaslan
 Tansılı Altay
 Tina Beşeri Sevim
 Turgay Akyar
 Tuğba Yazan
 Tuğrul Cömert
 Türker Özseri
 Türkü Özlüm Çelik
 Uyen Enni Le
 Victor Blasco
 Yaren Naz Tekingündüz
 Yasemin Soylu
 Yeşim Demiroğlu Karabulut
 Yıldız Oruklu
 Yılmaz Selçuk Bilgi
 Yonca Ünver
 Zehra Çayıç
 Zehra Tuncer
 Zeynep Üstün
 Zeynep Kara
 Zübeyir Türkoğlu
 Zümra Bulut

Dokuz Eylül University
 Muğla Sıtkı Koçman University
 İzmir Institute of Technology
 Yıldız Teknik University
 Yaşar University
 İzmir Institute of Technology
 Dokuz Eylül University
 Çukurova University
 Dokuz Eylül University
 Hacettepe University
 İzmir Institute of Technology
 Dokuz Eylül University
 Ankara Hacı Bayram Veli University
 Manisa Celal Bayar University
 İzmir Institute of Technology
 Dokuz Eylül University
 İzmir Institute of Technology
 İzmir Institute of Technology
 Hacettepe University
 İzmir Institute of Technology
 Rice University
 Hacettepe University
 İstanbul Bilgi University
 İzmir Institute of Technology
 Gebze Technical University
 İzmir Institute of Technology
 Dokuz Eylül University
 İzmir Institute of Technology
 İzmir Institute of Technology
 Middle East Technical University
 İzmir Institute of Technology
 Dicle University
 Bilkent University
 Boğaziçi University
 West Virginia University
 Dokuz Eylül University
 İzmir Institute of Technology
 Giresun University
 California Institute of Technology
 İzmir Institute of Technology
 İzmir Institute of Technology
 Amasya University
 İzmir Institute of Technology
 Dokuz Eylül University
 İzmir Institute of Technology
 Gazi University
 Dokuz Eylül University
 İzmir Institute of Technology